

DIFFERENTIATED SURGICAL TREATMENT OF PATIENTS WITH TRIGEMINAL NEURALGIA AND ASSESSMENT OF QUALITY OF LIFE**Mirzaev A.**

*Doctor of Medical Sciences, senior researcher, JSC "NMMC" MA "Kyzyl-Kum",
Deputy Director for Personnel and Administrative Issues, Republic of Uzbekistan
ORCID ID: <https://orcid.org/0000-0002-8216-54>*

Boymuradov Sh.,

*Doctor of Medical Sciences, Professor Vice-Rector for Academic Affairs,
Tashkent Medical Academy, Republic of Uzbekistan*

Abdullayev R.

*neurosurgeon, department of neurosurgery, multidisciplinary clinic,
Tashkent Medical Academy, Republic of Uzbekistan.
ORCID ID: <https://orcid.org/0009-0000-1986-136X>
<https://doi.org/10.5281/zenodo.13326972>*

Abstract

The article on the occurrence of trigeminal neuralgia, its surgical treatment, and the history of the development of other methods. Modern diagnostic methods, treatment tactics, determination of the effectiveness of treatment methods. Methods of assessing the quality of life the impact of a particular treatment method on the quality of life of patients. The analysis of the effectiveness of treatment methods is given. Application of questionnaires and comparative analysis of the results to determine the quality of life before and after the treatment of trigeminal neuralgia.

Keywords: neuralgia, trigeminal nerve, neurosurgery, surgical treatment, quality of life, questionnaires.

Relevance

Currently, we know two main causes of trigeminal neuralgia. They can be divided by the location of the origin of compression of the roots of the trigeminal nerve: intracranial and extracranial. Intracranial causes, when the compression of the root of the trigeminal nerve occurs at the brainstem. Extracranial compression occurs at the exits of the nerve roots through the opening in the skull: the supraorbital foramen is the first branch, the infraorbital foramen is the second branch, and the mandibular foramen is the third branch of the trigeminal nerve. The causes of compression can be inflammatory processes in these areas with extracranial lesions of the nerve roots of the trigeminal nerve [1,2,3,4,8,9,13].

At the present stage of development of medicine, a more valuable method of diagnosing trigeminal neuralgia is Magnetic resonance imaging (MRI) from 1.5 to 3.0 Tesla, by T2 (3D-FIESTA, DRIVE or CISS), time-of-flight magnetic resonance angiography (MRA) (3D-TOF), as well as contrast-enhanced weighted by T1 mode [1,2,3,4,8].

A more modern and very important method in diagnosing trigeminal neuralgia is electroneuromyography (ENMG), which determines the level of damage to the branches of the nerve, i.e. central or peripheral irritation. ENMG gives us the opportunity, taking into account the level of damage, to choose the tactics of surgical or conservative treatment of trigeminal neuralgia [1,2,3,4,8].

Currently, studying the literature, the authors note that MVD remains the method of choice if the patient is correctly selected, i.e. the established diagnosis is substantiated and proven by instrumental studies. As a result of the analysis, it was concluded that the MVD

provides the highest patient satisfaction rate in the long term with the lowest frequency of pain relapses in surgical treatment of trigeminal neuralgia [6,7,10,11,12,13,14,15].

In case of contraindications to other methods, manipulations on the peripheral fibers of the trigeminal nerve are performed at the patient's discretion. There are no protocols for their implementation; an individual and differentiated approach is always required [1,4,16].

Quality of life reflects the impact of the disease and treatment on the patient's well-being and characterizes his physical, emotional and social well-being, which changes under the influence of the disease or its treatment, therefore, scientists around the world have increasingly begun to study [5,10].

Purpose of the study: Improving the results of treatment of patients with trigeminal neuralgia by a differentiated approach taking into account the severity of the course and assessing the quality of life.

Material and methods

The study is based on the results of observations of 168 patients with trigeminal neuralgia, who were treated in the inpatient department of the Tashkent Medical Academy, a multidisciplinary clinic, and the neurosurgery department in the period from 2019 to 2023.

To establish a diagnosis and choose a treatment method, all patients underwent a comprehensive examination, including clinical, neurological and instrumental research methods.

The quality of life of patients was assessed using the questionnaires "Assessment of the quality of life in trigeminal neuralgia" and "Assessment of pain syn-

drome in trigeminal neuralgia" developed by the Republican Specialized Scientific and Practical Medical Center for Neurosurgery.

Results and discussion

The results of observations of 168 patients with trigeminal neuralgia, who were treated in the inpatient department of the Tashkent Medical Academy, a multidisciplinary clinic, and the neurosurgery department in the period from 2019 to 2023, were studied.

Before admission to the clinic, all patients received comprehensive conservative treatment, patients were treated by dentists, neurologists, maxillofacial surgeons.

All patients were divided into three groups according to treatment methods, severity of trigeminal neuralgia and somatic status.

The first group included 54 (32.2%) patients who underwent Gasser's node blockade, exeresis of the branches of the trigeminal nerve.

The second group consisted of 56 (33.3%) patients who underwent microvascular decompression surgery using a modified approach proposed by the Republican Specialized Scientific and Practical Medical Center for Neurosurgery for Trigeminal Neuralgia (TN).

The third group consisted of 58 (34.5%) patients who underwent the existing method of microvascular decompression of the trigeminal nerve root.

By age, patients are distributed according to the WHO classification, which provides for the allocation of age groups: young age 14-19 years; younger middle age 20-44 years; older middle age 45-59 years; elderly age 60-74 years; old age 75-89 years.

In our observations, patients were aged 24 to 79 years, taking this into account, the distribution of patients by age groups is as follows.

Distribution of patients by age and gender showed that among the patients, women predominated - 101 (60.2%), there were 67 men (39.8%), which is 1.5 times less than women. The majority of patients 69 (41.5%) were of older middle and old age, the maximum number of patients fell on the age group 60-74 years, 51 (30.4%), which is consistent with the data of world scientists.

When examining the somatic status, it was revealed that among 168 patients, 68 (40.5%) had somatic pathology, manifested as arterial hypertension in 52 (31.0%) patients, ischemic heart disease in 13 patients (7.8%). In 18 (11.2%) cases, diabetes mellitus was noted, in 4 (2.6%) patients, liver pathology, and in 1 (0.9%), renal failure. It should be noted that in one and the same patient, a combination of two or more somatic diseases was noted, this was especially characteristic of cardiovascular diseases.

Analysis of provoking factors based on the results of the study showed that almost all patients had: a provoking factor such as chewing and talking in more than 93% of cases, the second place is occupied by the factor of touch - 8% of observations, the rest are cold, brushing teeth, washing, shaving and opening the mouth in less than 5% of cases.

During examination of patients, the following

symptoms were identified as the main criteria of trigeminal neuralgia: unilateral paroxysmal facial pain, the presence of a trigger zone in the examined 168 patients, the localization of trigger zones gave the following results, with a total of 32 (19.3%) observations, among 32 patients with the presence of trigger points by localization, the nasolabial triangle ranks first 15 (48.5), the mental area ranks second 8 (24.3%), the frontal-zygomatic region ranks third 2 (9.1%), the remaining zones are the oral cavity, cheek, parotid region, Vale points, neck in the area of the spinous processes of the III-IV cervical vertebrae from 3% to 6% of cases.

We also checked for symptoms of hypertonicity of the masticatory muscles, autonomic disorders, hypesthesia of the innervation zones of individual branches or the entire half of the face. In this case, we were convinced that trigeminal neuralgia is characterized by the presence of short attacks (2-15 min), which can be repeated frequently (up to 30 times a day). The study of sensitivity disorders in the innervation zone of the trigeminal nerve branches yielded the following results.

In our study, out of 168 total, 90 (54%) patients had sensitivity disorders in the innervation zone of the trigeminal nerve branches: 52.2% of cases of hypesthesia, 40.2% of cases of paresthesia, 6.5% of cases of hyperesthesia, and 1 case of anesthesia as a result of previous surgical interventions. This case was observed among patients in the third group. As a result of multiple exeresis operations, sensitivity in the innervation zone was completely lost, and trophic changes in the facial muscles on the side of the affected branches occurred trigeminal nerve. This patient did not feel pain in the innervation zone of the trigeminal nerve after MVD, but sensitivity after destructive surgical interventions was not restored.

Vegetative disorders in our patients were observed in 17 (10.5%) cases, of 17 most of all 50.1% hyperemia, 33.3% lacrimation, 16.6% salivation if we consider by groups 55.6% in the third group, 27.7% in the second group and the least 16.7% in the first group, after treatment vegetative disorders were restored.

Analysis of the 168 patients examined, by the location of the affected branches of the trigeminal nerve showed the following results.

In 131 (78.4%) patients, trigeminal neuralgia was noted on the right, in 33 (19.8%) on the left and in 3 (1.8%) observations on both sides.

In our observations, the second (33.3%) and second + third branches (4.6%) were most often affected in the patients under study, and, as noted above, right-sided neuralgia was noted in most cases, which proves its typicality.

Thus, in our observations, 2 branches were most often affected simultaneously in 50.9% of patients. Damage to all three branches was observed in 31.9% and one branch in 17.2%. Most often, the pain syndrome was on the right in 78.4%, on the left in 19.8% and on both sides in 1.8% of cases. The data we obtained from the study of patients are consistent with the literature data of world authors.

Positive results among the 168 patients under study were noted in 164 (97.5%), without changes in 4 (2.5%). The best indicators were in the second and third

groups, all patients in these groups received positive results.

This proves the effectiveness of the surgical intervention method with a differentiated approach to the treatment of trigeminal neuralgia, taking into account the somatic status of the patient.

Analysis of the data obtained showed that 87.8% of patients in the second and third groups showed significant improvement, 12.2% showed improvement, no changes, in the first group 9.3% showed significant improvement, 83.4% showed improvement, and 4 (7.3%) patients showed no improvement.

In our studies, microvascular decompression of the trigeminal nerve root turned out to be a more effective surgical intervention method. Those operated on with microvascular decompression using a modified approach proposed by the Republican Specialized Scientific and Practical Medical Center for Neurosurgery for Trigeminal Neuralgia, the existing approach of microvascular decompression at the brainstem, since patients with significant improvements in the second and third groups amounted to 87.8% of the total number of 168 patients treated.

The use of the questionnaire "Assessment of the quality of life in trigeminal neuralgia" showed that the quality of life in all the studied groups was similar: that is, before the operation, the quality of life, indicators in all groups were low.

After the treatment, the quality of life, indicators immediately began to improve in patients with trigeminal neuralgia, this was especially noticeable in the second and third groups, and positive results were also shown by the treatment in patients of the first group.

The use of the questionnaire "Assessment of pain syndrome in trigeminal neuralgia" showed that the parameters of pain syndrome in the preoperative period in the studied patients in all groups differed from each other with a difference in points.

High indicators were recorded in the first group before the operation, which corresponded to a severe degree of severity of the pain syndrome, and after the treatment in all groups, a decrease in it to an average degree was observed in both the first and second groups, which was noted by the data obtained as a result of using the questionnaires.

In our studies, questionnaires from the Republican Specialized Scientific and Practical Medical Center of Neurosurgery were used to assess the quality of life and pain syndrome in patients with trigeminal neuralgia.

According to the questionnaire "Assessment of pain syndrome in trigeminal neuralgia": a sum of up to 10 points - mild severity of trigeminal neuralgia, 11-21 points - moderate severity of trigeminal neuralgia, 22-32 points - severe severity of trigeminal neuralgia. Assessment of pain syndrome in trigeminal neuralgia according to the questionnaire made it possible to determine in 168 patients mild pain syndrome in 40 (24.4%), moderate pain in 119 (71.3%), severe pain in 9 (5.2%), and in the postoperative period - a complete absence of pain was noted in 164 (97.3%) and in 5 (2.9%) - mild pain syndrome.

According to the questionnaire "Assessment of the quality of life in trigeminal neuralgia": a sum of up to

10 points is a slight deterioration in the quality of life, 11-21 points is a moderate deterioration in the quality of life, 22-33 points is a significant deterioration in the quality of life. Assessment of the quality of life according to the questionnaire in 168 patients with trigeminal neuralgia showed a significant deterioration in the quality of life in the preoperative period in 128 (76.5%) patients, and a deterioration in the quality of life in 40 (24.4%). And in the postoperative period, patients with a significant improvement in the quality of life amounted to 17 (10.4%) and with an improvement - 143 (85.4%), a slight improvement in the quality of life was noted in 7 (4.3%) patients.

Conclusions

1. Distribution of patients by age and gender showed that among the patients there were predominantly women - 101 (60.2%), there were 67 men (39.8%), which is 1.5 times less than women. Most patients 69 (41.5%) were of middle and old age, the maximum number of patients was in the age group of 60-74 years, 51 (30.4%), which is consistent with the data of world scientists.

2. When examining the somatic status, it was revealed that among 168 patients, 68 (40.5%) patients had somatic pathology, manifested in the form of arterial hypertension in 52 (31.0%) patients, ischemic heart disease in 13 patients (7.8%). In 18 (11.2%) observations, diabetes mellitus was noted, in 4 (2.6%) patients liver pathology and in 1 (0.9%) renal failure.

3. Positive results among the 168 patients studied were noted in 164 (97.5%), without changes in 4 (2.5%). The best indicators were in the second and third groups, all patients in these groups received positive results.

4. Evaluation of pain syndrome in trigeminal neuralgia using a questionnaire made it possible to determine in 168 patients mild pain syndrome in 40 (24.4%), moderate pain syndrome in 119 (71.3%), severe pain syndrome in 9 (5.2%), and in the postoperative period, a complete absence of pain was noted in 164 (97.3%) and in 5 (2.9%) - mild pain syndrome.

5. Assessment of the quality of life using a questionnaire in 168 patients with trigeminal neuralgia showed a significant deterioration in the quality of life in 128 (76.5%) patients in the preoperative period, and a deterioration in the quality of life in 40 (24.4%). And in the postoperative period, patients with a significant improvement in the quality of life amounted to 17 (10.4%) and with an improvement - 143 (85.4%), a slight improvement in the quality of life was noted in 7 (4.3%) patients.

References:

1. Alekseevets V.V., Likhachev S.A., Shanko Yu.G., Zmachinskaya O.L. / Modern approaches to the treatment of trigeminal neuralgia // Neurology and neurosurgery. Eastern Europe. - 2016. - No. 3. - P. 8-17. - Access mode: <https://rucont.ru/efd/525153> ISSN: 2226-0838eISSN: 2414-3588
2. Balyazina, E.V. / Comprehensive diagnostics of neurovascular conflict in patients with classical trigeminal neuralgia. // South-Russian Journal of Therapeutic Practice. - 2020. - 1 (1). - P. 70-77.

3. Balyazina, E.V., Balyazin V.A., Isakhanova T.A. et al. / Physical mechanism of formation of two types of neurovascular conflict in patients with classical trigeminal neuralgia. // *Neurological journal*. - 2017. - (4). - P. 190-197. - Access mode: doi: 10.18821/1560-9545-2017-22-4-190-197
4. Manvelov L.S., Tyurnikov V.M., Kadykov A.V. / Trigeminal neuralgia: epidemiology, etiology, pathomorphology, pathogenesis, clinical features, diagnostics. // *Zhurn. Vrach.* - 2013. - No. 4. - P. 6-8.
5. Novik A.A., Ionova T.I. / Guide to Quality of Life, Research in Medicine // edited by Yu. L. Shevchenko; Ministry of Health and Social Development of the Russian Federation, N. I. Pirogov National Medical and Surgical Center, Institute for Advanced Medical Studies of N. I. Pirogov National Medical and Surgical Center. - 3rd ed., revised and enlarged. - Moscow: Publishing House of the Russian Academy of Natural Sciences, 2012. - 527 p.: ill., port.; 21 cm; ISBN 978-5-94515-117-8 (In translation)
6. Rzaev D. A., Kulikova E. V., Moisak G. I., Voronina E. I., Ageeva T. A. / Teflon granuloma after microvascular decompression of the trigeminal nerve root in a patient with recurrent trigeminal neuralgia. // *Journal of Neurosurgery named after N.N. Burdenko*. 2016;80(2):78-83. <https://doi.org/10.17116/neiro201680278-83>
7. Guangyu Zhao, Xiaojun Sun, Zhen Zhang, Hongan Yang, Xiangrong Zheng, Bin Feng / Clinical efficacy of MVD combined with PSR in the treatment of primary trigeminal neuralgia // *Exp Ther Med*. 2020 Aug; 20(2):1582-1588. doi: 10.3892/etm.2020.8871. Epub 2020 Jun 10.
8. Latorre G, González-García N, García-Ull J, González-Oria C, Porta-Etessam J, Molina F J, Guerrero-Peral A I, Belvís R, Rodríguez R, Bescós A, Irimia P, Santos-Lasaosa S. / Diagnosis and treatment of trigeminal neuralgia: Consensus statement from the Spanish Society of Neurology's Headache Study Group // *Neurologia (Engl Ed)*. 2023 Apr 26; S2173-5808(23)00027-5. doi: 10.1016/j.nrleng.2023.04.005.
9. Matthew J. Allsop, Maureen Twiddy, Hilary Grant, Carolyn Czoski-Murray, Mark Mon-Williams, Faisal Mushtaq, Nick Phillips, Joanna M. Zakrzewska, Sue Pavitt / Diagnosis, medication, and surgical management for patients with trigeminal neuralgia: a qualitative study // *Acta Neurochir (Wien)*. 2015 Nov;157(11):1925-33. doi:10.1007/s00701-015-2515-4. Epub 2015 Sep 2.
10. Nicolas Jacques, Simon Karoutsos, Loïc Marais, Nathalie Nathan-Denizot / Quality of life after trigeminal nerve block in refractory trigeminal neuralgia: A retrospective cohort study and literature review // *J Int Med Res*. 2022 Oct;50(10):3000605221132027. doi: 10.1177/03000605221132027.
11. Ning Luo, Rongrong Li, Yiyi Wang, Yunfan Xia, Jing Sun, Linfang Zhao, Chao Sun, Jiemin Sun, Jianqiao Fang / Effectiveness and safety of acupuncture for treating trigeminal neuralgia associated anxiety and depression: A protocol for systematic review and meta-analysis // *Medicine (Baltimore)*. 2022 Jan 21;101(3):28631. doi: 10.1097/MD.00000000000028631.
12. Ruiz-Juretschke F, A J Vargas A.J., Gonzalez-Quarante L.H., Gil O L de Sagredo, Montalvo A, / C Fernandez-Carballal Microsurgical treatment of trigeminal neuralgia in patients older than 70 years: An efficacy and safety study // *Neurologia*. 2017 Sep;32(7):424-430. doi: 10.1016/j.nrl.2016.01.007. Epub 2016 Mar 9.
13. Wataru Yoshizaki, Yoshiki Fujikawa, Sadaharu Torikoshi, Toshiro Katayama, Koichi Iwasaki, Hiroki Toda / Effects of microvascular decompression on quality-of-life in trigeminal neuralgia patients aged 70 years and older // *Surg Neurol Int*. 2023 Feb 3;14:41. doi: 10.25259/SNI_997_2022. eCollection 2023.
14. Yejiào Luo, Mingjie He, Chenjun Li, Hongya Yang / A research on quality of life, score (QOLS) of patients with trigeminal neuralgia (TN) // *J Infect Public Health*. 2019 Sep-Oct;12(5):690-694. doi: 10.1016/j.jiph.2019.03.011. Epub 2019 May 14.
15. Yusuke Otsu, Sosho Kajiwarra, Aya Hashimoto, Kiyohiko Sakata, Tetsuya Negoto, Yu Hasegawa, Hideo Nakamura, Masaru Hirohata, Motohiro Morioka / Effects of Microvascular Decompression on Pain Relief and Quality of Life in Late Elderly Patients with Trigeminal Neuralgia // *Neurol Med (Tokyo)*. 2023 Jun 15;63(6):236-242. doi: 10.2176/jns-nmc.2022-0288. Epub 2023 Apr 6.